

Undetermined coefficients in constant coefficient ODEs

Prof. Joyner¹

The method of undetermined coefficients [U] can be used to solve the following type of problem.

PROBLEM: Solve

$$ay'' + by' + cy = f(t), \quad (1)$$

where $a \neq 0$, b and c are constants. (Even the case $a = 0$ can be handled similarly, though some of the discussion below might need to be slightly modified.) Where we must assume that $f(t)$ is of a special form.

More-or-less equivalent is the method of annihilating operators [A] (they solve the same class of DEs), but that method will be discussed separately.

For the moment, let us assume $f(t)$ has the form $a_1 \cdot p(t) \cdot e^{a_2 t} \cdot \cos(a_3 t)$, or $a_1 \cdot p(t) \cdot e^{a_2 t} \cdot \sin(a_3 t)$, where a_1, a_2, a_3 are constants and $p(t)$ is a polynomial.

Solution:

- Find the “homogeneous solution” y_h to $ay'' + by' + cy = 0$, $y_h = c_1 y_1 + c_2 y_2$. Here y_1 and y_2 are determined as follows: let r_1 and r_2 denote the roots of the characteristic polynomial $aD^2 + bD + c = 0$.
 - $r_1 \neq r_2$ real: set $y_1 = e^{r_1 t}$, $y_2 = e^{r_2 t}$.
 - $r_1 = r_2$ real: if $r = r_1 = r_2$ then set $y_1 = e^{rt}$, $y_2 = te^{rt}$.
 - r_1, r_2 complex: if $r_1 = \alpha + i\beta$, $r_2 = \alpha - i\beta$, where α and β are real, then set $y_1 = e^{\alpha t} \cos(\beta t)$, $y_2 = e^{\alpha t} \sin(\beta t)$.
- Compute $f(t)$, $f'(t)$, $f''(t)$, Write down the list of all the different terms which arise (via the product rule), ignoring constant factors, plus signs, and minus signs:

¹These notes licensed under Attribution-ShareAlike Creative Commons license, <http://creativecommons.org/about/licenses/meet-the-licenses>. Created Feb 2007; last revised Nov 2, 2007.

$$f_1(t), f_2(t), \dots, f_k(t).$$

If any one of these agrees with y_1 or y_2 then multiply them all by t . (If, after this, any of them *still* agrees with y_1 or y_2 then multiply them all again by t .)


- Let y_p be a linear combination of these functions (your “guess”):

$$y_p = A_1 f_1(t) + \dots + A_k f_k(t).$$

This is called the **general form of the particular solution**. The A_i 's are called **undetermined coefficients**.

- Plug y_p into (1) and solve for A_1, \dots, A_k .
- Let $y = y_h + y_p = y_p + c_1 y_1 + c_2 y_2$. This is the **general solution** to (1). If there are any initial conditions for (1), solve for then c_1, c_2 now.

Diagrammatically:



Examples

Example 1 Solve

$$y'' - y = \cos(2x).$$

- The characteristic polynomial is $r^2 - 1 = 0$, which has ± 1 for roots. The “homogeneous solution” is therefore $y_h = c_1 e^x + c_2 e^{-x}$.
- We compute $f(x) = \cos(2x)$, $f'(x) = -2 \sin(2x)$, $f''(x) = -4 \cos(2x)$, They are all linear combinations of

$$f_1(x) = \cos(2x), \quad f_2(x) = \sin(2x).$$

None of these agrees with $y_1 = e^x$ or $y_2 = e^{-x}$, so we do not multiply by x .

- Let y_p be a linear combination of these functions:

$$y_p = A_1 \cos(2x) + A_2 \sin(2x).$$

- You can compute both sides of $y_p'' - y_p = \cos(2x)$:

$$(-4A_1 \cos(2x) - 4A_2 \sin(2x)) - (A_1 \cos(2x) + A_2 \sin(2x)) = \cos(2x).$$

Equating the coefficients of $\cos(2x)$, $\sin(2x)$ on both sides gives 2 equations in 2 unknowns: $-5A_1 = 1$ and $-5A_2 = 0$. Solving, we get $A_1 = -1/5$ and $A_2 = 0$.

- The general solution: $y = y_h + y_p = c_1 e^x + c_2 e^{-x} - \frac{1}{5} \cos(2x)$.

Example 2 Solve

$$y'' - y = x \cos(2x).$$

- The characteristic polynomial is $r^2 - 1 = 0$, which has ± 1 for roots. The “homogeneous solution” is therefore $y_h = c_1 e^x + c_2 e^{-x}$.
- We compute $f(x) = x \cos(2x)$, $f'(x) = \cos(2x) - 2x \sin(2x)$, $f''(x) = -2 \sin(2x) - 2 \sin(2x) - 2x \cos(2x)$, They are all linear combinations of

$$f_1(x) = \cos(2x), \quad f_2(x) = \sin(2x), \quad f_3(x) = x \cos(2x), \quad f_4(x) = x \sin(2x).$$

None of these agrees with $y_1 = e^x$ or $y_2 = e^{-x}$, so we do not multiply by x .

- Let y_p be a linear combination of these functions:

$$y_p = A_1 \cos(2x) + A_2 \sin(2x) + A_3 x \cos(2x) + A_4 x \sin(2x).$$

- In principle, you can compute both sides of $y_p'' - y_p = x \cos(2x)$ and solve for the A_i 's. (Equate coefficients of $x \cos(2x)$ on both sides, equate coefficients of $\cos(2x)$ on both sides, equate coefficients of $x \sin(2x)$ on both sides, and equate coefficients of $\sin(2x)$ on both sides. This gives 4 equations in 4 unknowns, which can be solved.) You will not be asked to solve for the A_i 's for a problem this hard.

Example 3 Solve

$$y'' + 4y = x \cos(2x).$$

- The characteristic polynomial is $r^2 + 4 = 0$, which has $\pm 2i$ for roots. The "homogeneous solution" is therefore $y_h = c_1 \cos(2x) + c_2 \sin(2x)$.
- We compute $f(x) = x \cos(2x)$, $f'(x) = \cos(2x) - 2x \sin(2x)$, $f''(x) = -2 \sin(2x) - 2 \sin(2x) - 2x \cos(2x)$, They are all linear combinations of

$$f_1(x) = \cos(2x), \quad f_2(x) = \sin(2x), \quad f_3(x) = x \cos(2x), \quad f_4(x) = x \sin(2x).$$

Two of these agree with $y_1 = \cos(2x)$ or $y_2 = \sin(2x)$, so we do multiply by x :

$$f_1(x) = x \cos(2x), \quad f_2(x) = x \sin(2x), \quad f_3(x) = x^2 \cos(2x), \quad f_4(x) = x^2 \sin(2x).$$

- Let y_p be a linear combination of these functions:

$$y_p = A_1 x \cos(2x) + A_2 x \sin(2x) + A_3 x^2 \cos(2x) + A_4 x^2 \sin(2x).$$

- In principle, you can compute both sides of $y_p'' + 4y_p = x \cos(2x)$ and solve for the A_i 's. You will not be asked to solve for the A_i 's for a problem this hard.

More generally, suppose that you want to solve $ay'' + by' + cy = f(x)$, where $f(x)$ is a sum of functions of the above form. In other words, $f(x) = f_1(x) + f_2(x) + \dots + f_k(x)$, where each $f_j(x)$ is of the form $c \cdot p(x) \cdot e^{ax} \cdot \cos(bx)$, or $c \cdot p(x) \cdot e^{ax} \cdot \sin(bx)$, where a, b, c are constants and $p(x)$ is a polynomial. You can proceed in either one of the following ways.

1. Split up the problem by solving each of the k problems $ay'' + by' + cy = f_j(x)$, $1 \leq j \leq k$, obtaining the solution $y = y_j(x)$, say. The solution to $ay'' + by' + cy = f(x)$ is then $y = y_1 + y_2 + \dots + y_k$ (the superposition principle).
2. Proceed as in the examples above but with the following slight revision:
 - Find the “homogeneous solution” y_h to $ay'' + by' + cy = 0$, $y_h = c_1y_1 + c_2y_2$.
 - Compute $f(x)$, $f'(x)$, $f''(x)$, Write down the list of all the different terms which arise, ignoring constant factors, plus signs, and minus signs:

$$t_1(x), t_2(x), \dots, t_k(x).$$

- Group these terms into their *families*. Each family is determined from its parent(s) - which are the terms in $f(x) = f_1(x) + f_2(x) + \dots + f_k(x)$ which they arose from by differentiation. For example, if $f(x) = x \cos(2x) + e^{-x} \sin(x) + \sin(2x)$ then the terms you get from differentiating and ignoring constants, plus signs and minus signs, are

$$x \cos(2x), x \sin(2x), \cos(2x), \sin(2x), \quad (\text{from } x \cos(2x)),$$

$$e^{-x} \sin(x), e^{-x} \cos(x), \quad (\text{from } e^{-x} \sin(x)),$$

and

$$\sin(2x), \cos(2x), \quad (\text{from } \sin(2x)).$$

The first group absorbs the last group, since you can only count the *different* terms. Therefore, there are only two families in this example: $\{x \cos(2x), x \sin(2x), \cos(2x), \sin(2x)\}$ is a “family” (with “parent” $x \cos(2x)$ and the other terms as its “children”) and $\{e^{-x} \sin(x), e^{-x} \cos(x)\}$ is a “family” (with “parent” $e^{-x} \sin(x)$ and the other term as its “child”).

If any one of these terms agrees with y_1 or y_2 then multiply the *entire family* by x . In other words, if any child or parent is “bad” then the entire family is “bad”. (If, after this, any of them *still* agrees with y_1 or y_2 then multiply them all again by x .)

- Let y_p be a linear combination of these functions (your “guess”):

$$y_p = A_1 t_1(x) + \dots + A_k t_k(x).$$

This is called the **general form of the particular solution**.
The A_i 's are called **undetermined coefficients**.

- Plug y_p into (1) and solve for A_1, \dots, A_k .
- Let $y = y_h + y_p = y_p + c_1 y_1 + c_2 y_2$. This is the **general solution** to (1). If there are any initial conditions for (1), solve for then c_1, c_2 last - after the undetermined coefficients.

Example 4 Solve

$$y''' - y'' - y' + y = 12xe^x.$$

We use SAGE for this.

SAGE

```
sage: x = var("x")
sage: y = function("y",x)
sage: R.<D> = PolynomialRing(QQ, "D")
sage: f = D^3 - D^2 - D + 1
sage: f.factor()
(D + 1) * (D - 1)^2
sage: f.roots()
[(-1, 1), (1, 2)]
```

So the roots of the characteristic polynomial are 1, 1, -1, which means that the homogeneous part of the solution is

$$y_h = c_1 e^x + c_2 x e^x + c_3 e^{-x}.$$

SAGE

```
sage: de = lambda y: diff(y,x,3) - diff(y,x,2) - diff(y,x,1) + y
sage: c1 = var("c1"); c2 = var("c2"); c3 = var("c3")
sage: yh = c1*e^x + c2*x*e^x + c3*e^(-x)
```

```
sage: de(yh)
0
sage: de(x^3*e^x-(3/2)*x^2*e^x)
12*x*e^x
```

This just confirmed that y_h solves $y''' - y'' - y' + 1 = 0$. Using the derivatives of $F(x) = 12xe^x$, we generate the general form of the particular:

SAGE

```
sage: F = 12*x*e^x
sage: diff(F,x,1); diff(F,x,2); diff(F,x,3)
12*x*e^x + 12*e^x
12*x*e^x + 24*e^x
12*x*e^x + 36*e^x
sage: A1 = var("A1"); A2 = var("A2")
sage: yp = A1*x^2*e^x + A2*x^3*e^x
```

Now plug this into the DE and compare coefficients of like terms to solve for the undertermined coefficients:

SAGE

```
sage: de(yp)
12*x*e^x*A2 + 6*e^x*A2 + 4*e^x*A1
sage: solve([12*A2 == 12, 6*A2+4*A1 == 0],A1,A2)
[[A1 == -3/2, A2 == 1]]
```

Finally, lets check if this is correct:

SAGE

```
sage: y = yh + (-3/2)*x^2*e^x + (1)*x^3*e^x
sage: de(y)
12*x*e^x
```

Exercise: Using SAGE, solve

$$y''' - y'' + y' - y = 12xe^x.$$

(Hint: You may need to replace `sage: R.<D> = PolynomialRing(QQ, "D")` by `sage: R.<D> = PolynomialRing(CC, "D").`)

References

[A] Wikipedia entry for the annihilator method:
http://en.wikipedia.org/wiki/Annihilator_method

[U] General wikipedia introduction to undetermined coefficients:
http://en.wikipedia.org/wiki/Method_of_undetermined_coefficients